



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.


We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

Solution by G. I. HOPKINS, A. M., Professor of Mathematics and Astronomy, Manchester (N. H.) High School, and MISS IDA M. SCHOTTENFELS, A. M., New York, N. Y.

Let  $A, B, C, D, E, F, G, H,$  and  $K$  be the points. It is evident that  $A, C, K,$  and  $G$  will be the vertices of a parallelogram. Let  $BH$  be a median of this parallelogram, and  $E$  any point in the median except the center. Then the broken line  $ANPRK$  will fulfill the conditions of the problem. This course fails if the middle row and middle column bisect each other. If the row  $DEF$  is not parallel to  $GK$ , then three lines, or a broken line of three segments will fulfill the conditions of the problem.



G. I. HOPKINS.

If  $BH$  and  $DF$  are not medians, take the course  $KER, RP, PN, NB$ . If  $BH$  and  $DF$  are medians, take the course  $KEA, AN, NP, PD$ ; or  $KEA, AP, PN, NB$ .

IDA M. SCHOTTENFELS.

Also solved by the Proposer.

296 (Incorrectly numbered 294). Proposed by JOHN JAMES QUINN, Ph. D., Scottsdale, Pa.

a) Suppose an indefinite line be pivoted at the end of a revolving radius whose center is the origin; and the initial position of the radius is coincident with the  $X$ -axis and the pivoted line perpendicular to it. As the radius revolves through equal amounts of arc the line moves to the right over corresponding equal intercepts on the  $X$ -axis. What is the equation of the locus of a point on the line whose distance from the end of the radius is equal to a diameter?

b) Show how the locus can be applied to the multisection of an angle.

c) Suppose the diameter be laid off in both directions.

No solution of this problem has been received.

297 (Incorrectly numbered 295). Proposed by S. F. NORRIS, Professor of Mathematics, Baltimore City College, Md.

One side and the opposite angle of a triangle are fixed. Find the locus of the center of the inscribed circle. Solve by methods of analytic geometry.

I. Solution by C. N. SCHMALL, A. B., 89 Columbia Street, New York City.

This problem can be solved more easily and more neatly by Euclidean Geometry. Thus, referring to figure, we have,


$$\angle DCO = \angle DCB + \angle BCO = \angle DCB + \frac{1}{2} \angle BCA,$$

$$\angle DOC = \angle AOF = \angle OAC + \angle OCA;$$

but  $\angle OAC = \angle DCB$  (since arc  $DC = DB$ ), and  $\angle OCA = \frac{1}{2} \angle BCA$ . Hence  $\angle DCO = \angle DOC$ .

Therefore  $DC = DO$ .

Hence, keeping  $BC$  constant and vertex  $A$  always on the arc  $BAC$  (making opposite angle constant) the locus of center  $O$  of the inscribed circle is a circle whose center is  $D$  and radius  $DC$ .



II. Solution by G. W. GREENWOOD, A. M., Dunbar, Pa.

Taking as  $x$ -axis the fixed side  $AB$  ( $=2a$ ) and its mid-point as origin, the in-center, since it lies on the bisectors of the angles  $A$  and  $B$ , must satisfy the equations  $y=m_1(x-a)$ ,  $y=m_2(x+a)$ , where

$$m_1=\tan\left(\pi-\frac{A}{2}\right)=-\tan\frac{A}{2}, \quad m_2=\tan\frac{B}{2}.$$

The opposite angle being constant,  $A+B$  is constant. Hence

$$\tan\left(\frac{A}{2}+\frac{B}{2}\right)=\frac{-m_1+m_2}{1+m_1m_2}=\text{a constant}=c, \text{ say.}$$

Hence the in-center satisfies the equation  $c(x^2+y^2-a^2)+2ay=0$ .

Also solved by J. Scheffer, William Hoover, and L. E. Newcomb.

298 (Incorrectly numbered 296). Proposed by J. J. QUINN, Ph. D., Scottdale, Pa.

Given  $AB=BC$  perpendicular to each other, and  $E$  and  $M$  their mid-points, respectively. On  $AB$  describe a semi-circle, and draw  $CE$  to meet the circumference in  $D$ . Draw  $DM$  cutting  $AB$  in  $F$ . In what ratio is  $AB$  divided by the point  $F$ ?

Solution by C. N. SCHMALL, A. B., 89 Columbia Street, New York; L. E. NEWCOMB, Los Gatos, California; and A. H. HOLMES, Brunswick, Maine.

From the figure, constructed as described in the problem, we have

$$FE:ED=NM:ND. \quad \text{But } NM=\frac{1}{2}EB, \text{ and}$$

$$ND=CD-CN=CD-\frac{1}{2}CE...(1).$$

$$\therefore FE:ED=\frac{1}{2}EB:CD-\frac{1}{2}CE...(2).$$

Also  $AB^2=CD.CD'$ , where  $D'$  is the second intersection of the secant,  $CE$ , with the circle. Now let  $FA=x$ ,  $EB=r$ ,  $CD=a$ . Then  $FE=x+r$ ,  $CE=a+r$ . Substituting these values in (1) and (2), we have

$$x+r:r=\frac{1}{2}r:a-\frac{1}{2}(a+r)...(3),$$

$$(2r)^2=a(a+2r)...(4).$$

From (4),  $a=r(\pm\sqrt{5}-1)$ , and from (3),

$$a=r+\frac{r^2}{x+r}. \quad \therefore r(\pm\sqrt{4}-1)=r+\frac{r^2}{x+r}.$$

Whence  $x=(\pm\sqrt{5}+1)r$ .  $\therefore FA/AB=\frac{1}{2}(\pm\sqrt{5}+1)$ .

Also solved by G. W. Greenwood and J. Scheffer.

